

Sea Duck Joint Venture Final Report

Project Title: Evaluating stable hydrogen isotopes for identifying breeding areas of harvested sea ducks (F21AC00061-00)

Principal Investigators: Dr. David Nelson (University of Maryland Center for Environmental Science, dnelson@umces.edu); Dr. Hannah Vander Zanden (University of Florida, hvz@ufl.edu); Dr. Todd Katzner (United States Geological Survey, tkatzner@usgs.gov)

Partners: None beyond SDJV.

Project Description: This project represents a pilot study aimed at assessing the ability of feather stable hydrogen isotope ($\delta^2\text{H}$) data to delineate the most probable breeding locations of juvenile long-tailed ducks and surf scoters harvested in North America. To accomplish our objectives, SDJV provided us with feathers from 120 harvested individuals in the SDJV's feather archive (n= 60 juvenile long-tailed ducks and n = 60 juvenile surf scoters) spread across the Atlantic, Mississippi, and Pacific flyways. For 90% (n=108) of these individuals we performed $\delta^2\text{H}$ analysis on a single feather, whereas for 10% of them (n=12) we assessed intra-feather and intra-individual variation in isotope values by measuring subsamples from the proximal, middle and distal ends of 3 feathers per individual. Following an initial evaluation of the $\delta^2\text{H}$ data that we obtained from these feathers (see Results below), we suspected that some individuals may have been provisioned with marine resources that would make them unsuitable for geolocation purposes using $\delta^2\text{H}$ data. Therefore, to identify such individuals we performed carbon and nitrogen isotope ($\delta^{13}\text{C}$ and $\delta^{15}\text{N}$) analyses of the same feather samples. Results of the pilot study funded under agreement F21AC00061-00 identified that feather samples of several surf scoters and long-tailed ducks showed potential marine influence in their diet prior to fledging. However, this is contrary to the current understanding that these species (especially surf scoters) typically rear their young in freshwater/terrestrial environments. The SDJV Continental Technical Team then requested that we then conduct additional analysis of feathers from live-captured birds of known age (provided by SDJV) to investigate whether previously sampled birds were misidentified as hatch year birds.


Project Objectives: (1) Quantify intra-feather and intra-individual variation in feather $\delta^2\text{H}$ values, (2) Use $\delta^2\text{H}$ values to create a likelihood-of-origin maps, (3) Use likelihood-of-origin maps to identify regions where juveniles fledged and grew their feathers, (4) Evaluate the need to improve geo-location through isotopic analysis of feathers from reference individuals with known breeding locations, and (5) If results indicate that $\delta^2\text{H}$ values are useful for estimating breeding locations, then provide a preliminary evaluation of hypotheses about variation in the location of breeding grounds of these species between flyways. The amendment to the project added the following 6th objective: Test whether feathers from known-aged scoters and long-tailed ducks also show marine influence in their diet via $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$ analysis.

Results: For the 12 birds for which we sampled from multiple feathers and from multiple locations within feathers, intra- and inter-feather in $\delta^2\text{H}$, $\delta^{13}\text{C}$, and $\delta^{15}\text{N}$ values was small relative to the among-bird variation. The mean (range) standard deviation of multiple analyses per feather was 6.6 ‰ (2.1-17.5 ‰), 0.9 ‰ (0.1-3.5 ‰), and 0.5 ‰ (0.1-1.2 ‰), respectively, across all 12 birds, indicating that intra-variation in $\delta^2\text{H}$, $\delta^{13}\text{C}$, and $\delta^{15}\text{N}$ values was relatively small. The mean standard deviation across all 12 birds of all feathers from each bird was 7.3, 0.9, and 0.5 ‰ for $\delta^2\text{H}$, $\delta^{13}\text{C}$, and $\delta^{15}\text{N}$, respectively, indicating that, for a given bird, inter-feather isotopic variation was similar in magnitude to intra-feather isotopic variation. These patterns are consistent with our expectation that intra- and inter-feather isotopic variation be relatively small since juvenile sea ducks are believed to form all of their feathers in one general location

within a relatively short period of time. Therefore, isotopic data from one feather of a juvenile should be representative of isotopic values of that individual.

The results from these 12 birds also revealed a much larger range of variation in $\delta^2\text{H}$ values than would be expected for individuals that grew their feathers in the northern high latitudes where these species are born and grow their feathers (Fig. 1). Specifically, the highly positive $\delta^2\text{H}$ values of some individuals would suggest that they fledged far south of their expected range, assuming they were feeding in terrestrial and/or freshwater food webs when their feathers were formed. However, an alternative possibility is that these individuals may have been eating from coastal/marine food webs during the period of feather growth. To assess this possibility, we measured $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$ values for these 12 individuals, since prior studies indicate that $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$ values tend to be greater than approximately -20 and 11 ‰, respectively, for birds eating from coastal/marine food webs relative to those eating from terrestrial and/or freshwater food webs (Chamberlain et al. 2005, Yerkes et al. 2008, Newsome et al. 2010). Four individuals (two of each species) had $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$ values suggestive of marine inputs (data not shown), and thus the $\delta^2\text{H}$ values of these birds are likely ill-suited for geolocation purposes.

We also performed $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$ analyses of the single feathers from each of the 108 other birds that were part of our study. Overall, 8/60 long-tailed ducks (4/30 from the Atlantic flyway and 4/30 from the Mississippi flyway) and 12/60 surf scoters (10/19 from Pacific flyway, 0/11 from the Mississippi flyway, and 2/30 from the Atlantic flyway) had $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$ values suggestive of a marine influence in their diets (Fig. 2). Therefore, most individuals of each species appear to have hatched and fledged in terrestrial/freshwater habitat. The existence of some juveniles with possible coastal/marine diets is generally contrary to the current understanding that these species (especially surf scoters) are hatched and raised in terrestrial and/or freshwater environments. One explanation may be that the age of those hunter-killed birds was misidentified at wing bees (i.e., instead of being hatch-year individuals, perhaps they were older birds). Alternatively, the apparent marine signatures may be related to nutrient input from anadromous Pacific salmon, which migrate from the ocean hundreds of miles inland, mainly in Alaska and northwestern Canada, and die, thereby delivering marine-derived nutrients to potential sea duck breeding areas (Schindler et al. 2003, Reisinger et al. 2013). This would help explain why there was a higher proportion of apparent marine signatures in some Pacific surf scoters. To help distinguish among these possibilities, in spring 2022 the SDJV Continental Technical Team provided us with additional feathers from lived-captured long-tailed ducks (n=44 from




Atlantic flyway) and surf scoters (n=30 from Pacific flyway and n=30 from Atlantic flyway) whose age (i.e., hatch year) was definitively known; those analyses are in progress.

Several lines of evidence indicate that $\delta^2\text{H}$ values are a promising approach for delineating the breeding locations of juvenile long-tailed ducks and surf scoters harvested in North America. First, $\delta^2\text{H}$ values for the hatch-year long-tailed ducks and surf scoters eating on freshwater/terrestrial food webs (n=52 and n=48, respectively) fall within the ranges expected for individuals from throughout their geographic ranges (Fig. 1). Second, the surf scoters tended to have more positive $\delta^2\text{H}$ values (mean = -116‰) than long-tailed ducks (mean = -131‰), consistent with surf scoters having a relatively more southern breeding distribution (Fig. 1). Third, the locations at which the birds were killed were south of their $\delta^2\text{H}$ -inferred geographic origins (data not shown), as expected. Fourth, the

$\delta^2\text{H}$ values allowed us to identify the 20% of the species range (on average) from which each individual originated, which illustrates the precision at which we can infer fledgling locations using this approach. These results, along with low intra- and inter-feather isotopic variation, indicate that $\delta^2\text{H}$ values are useful for constraining the breeding locations of juvenile long-tailed ducks and surf scoters harvested in North America, even without feathers from individuals of these species with known breeding locations.

Our results indicate that juvenile long-tailed ducks likely provisioned from terrestrial/freshwater food webs had a large range of $\delta^2\text{H}$ values (Fig. 1), indicating origins spanning from northern Quebec to Nunavut (Fig. 3). Mean $\delta^2\text{H}$ values from the Atlantic and Mississippi flyways were similar (-131 ‰; data not shown), which suggests that individuals killed in these flyways have overlapping geographic origins. This result stands in contrast to the expectation from telemetry data that individuals from the Mississippi flyway would breed to the south and west of those from the Atlantic flyway (https://seaduckjv.org/wp-content/uploads/2014/12/AGLSDMS-Progress-Report-June2015_web.pdf). However, in our existing long-tailed duck dataset (1) sample sizes are relatively low (n=30 birds/flyway), (2) there are only two individuals that were killed north of Massachusetts (i.e., sample representation is poor in the maritime proveniences of Canada), (3) half of the samples from the Mississippi flyway are missing information on the location and date of harvest and (4) there are no samples from birds killed in the Pacific flyway, including Alaska and British Columbia.

For juvenile surf scoters, our results indicate that individuals provisioned from terrestrial/freshwater food webs also had a large range of $\delta^2\text{H}$ values (Fig. 1), indicating origins spanning from central Quebec to northern western Canada and Alaska (Fig. 3). Individuals from the Atlantic flyway tended to have higher $\delta^2\text{H}$ values (mean = -109 ‰) than those from the Mississippi (mean = -133 ‰) and Pacific (mean = -125 ‰) flyways (data not shown), which suggests a more southerly geographic origins in eastern Canada for individuals shot in the Atlantic flyway than for individuals from the latter two flyways that likely originate from northwestern Canada and Alaska. However, in the existing dataset



(1) sample sizes are relatively low and (2) there are no hunter-killed samples in the Pacific flyway from areas north of Washington state, which suggests that the existing samples from the Pacific flyway are likely not representative of that flyway.

We incorporated feather $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$ data into our pilot project once we suspected that some individuals may have been provisioned on coastal/marine food webs, as this approach has been used in prior studies (Chamberlain et al. 2005, Yerkes et al. 2008, Newsome et al. 2010) and we could readily perform these analyses in our lab. However, it is possible for birds provisioned on freshwater systems to have $\delta^{13}\text{C}$ values greater than -20‰ as a result of several factors, such as variation in the concentrations and $\delta^{13}\text{C}$ values of dissolved inorganic carbon (O'Leary 1988, Bade et al. 2004, Campeau et al. 2017); it is also possible for birds provisioned on terrestrial/freshwater systems to have $\delta^{15}\text{N}$ values greater than 11‰ because of trophic fractionation of nitrogen isotopes (Hobson 1999, Fry 2006). Sulfur isotopes ($\delta^{34}\text{S}$) would provide additional help for discriminating power between sea ducks provisioned in terrestrial/freshwater and coastal/marine ecosystems. The $\delta^{34}\text{S}$ values of marine-derived sulfur are generally more positive values than terrestrial sources and $\delta^{34}\text{S}$ values is not influenced by trophic fractionation of isotopes (Hache et al. 2014). Prior studies have used a threshold of $\delta^{34}\text{S}$ value of $\geq 11\text{‰}$ to identify feathers of coastal/marine provenance. Such data can improve spatial assignments based on $\delta^2\text{H}$, as well as help to identify the importance of coastal habitats as potential fledgling areas more definitively (Hobson 1999, Lott et al. 2003, Hobson and Kardynal 2016). As with $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$, there can be cases where birds in terrestrial/freshwater systems with unique hydrogeochemistry have $\delta^{34}\text{S}$ values of $\geq 11\text{‰}$ (Hitchon and Krouse 1972), but it would be unlikely for the combination of $\delta^{13}\text{C}$, $\delta^{15}\text{N}$, and $\delta^{34}\text{S}$ to incorrectly classify individuals provisioned on terrestrial/freshwater vs. coastal/marine resources.

For the live-captured juvenile surf scoters and long-tailed duck samples that were provided to us we found that that only 1 of 48 (or 2%) of surf scoters, including 1 of 30 from the Atlantic flyway and 0 of 28 from the Pacific fly way seem to have a marine influence (Fig. 4). This result contrasts to the wingbee data, which showed that 20% of surf scoters had a marine influence, including over half of those from the Pacific flyway. Thus, it seems probable that some of the wingbee surf scoters samples that were identified as juveniles were older birds that were eating on marine food webs. In contrast, for the long-tailed ducks, we observed that 5 of 49 or 10% of the live-captured birds from the Atlantic flyway had a marine influence, which is similar to the percent we observed in the wingbee data. Thus, there does seem to be a small proportion of long-tailed duck nestlings that are provisioned using marine resources, at least






Figure 3. Composite likelihood-of-origin maps for 60 long-tailed ducks (top) and 60 surf scoters (bottom). Dark blue/purple colors indicate areas where more individuals have shared geographic origin than the lighter green/yellow colors.

based on the $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$ data. Furthermore, similar to the wingbee data, the non-marine surf scoter samples tend to plot farther from the terrestrial/marine threshold than do the long-tailed ducks, which may be related to differences in their habitat use and diet. As expected, live-captured individuals with the more positive $\delta^2\text{H}$ values are typically those with marine inputs to diets, with the exception of one long-tailed duck with $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$ values suggestive of marine diet that had a relative negative $\delta^2\text{H}$ value (Fig. 5).

Project Status: We completed the project objectives. We believe that sulfur isotopes ($\delta^{34}\text{S}$) may provide additional help for discriminating power between sea ducks

provisioned in terrestrial/freshwater and coastal/marine ecosystems. We have recently received notice that a new proposal we submitted to SDJV will receive funding, which will allow us to expand these existing results and fund a graduate student researcher to (1) more rigorously assess variation in the location of breeding habitat across flyways and (2) better evaluate the potential of coastal/marine habitat as fledgling areas.



References

- Bade, D. L., S. R. Carpenter, J. J. Cole, P. C. Hanson, and R. H. Hesslein. 2004. Controls of delta $\delta^{13}\text{C}$ -DIC in lakes: Geochemistry, lake metabolism, and morphometry. *Limnology and Oceanography* **49**:1160-1172.
- Campeau, A., M. B. Wallin, R. Giesler, S. Lofgren, C. M. Morth, S. Schiff, J. J. Venkiteswaran, and K. Bishop. 2017. Multiple sources and sinks of dissolved inorganic carbon across Swedish streams, refocusing the lens of stable C isotopes. *Scientific Reports* **7**.
- Chamberlain, C. P., J. R. Waldbauer, K. Fox-Dobbs, S. D. Newsome, P. L. Koch, D. R. Smith, M. E. Church, S. D. Chamberlain, K. J. Sorenson, and R. Risebrough. 2005. Pleistocene to recent dietary shifts in California condors. *Proceedings of the National Academy of Sciences of the United States of America* **102**:16707-16711.
- Fry, B. 2006. *Stable Isotope Ecology*. Springer-Verlag, New York.
- Hache, S., K. A. Hobson, E. M. Bayne, S. L. Van Wilgenburg, and M. A. Villard. 2014. Tracking natal dispersal in a coastal population of a migratory songbird using feather stable isotope ($\delta^2\text{H}$, $\delta^{34}\text{S}$) tracers. *Plos One* **9**.
- Hitchon, B., and H. R. Krouse. 1972. Hydrogeochemistry of surface waters of Mackenzie River drainage basin, Canada- III. Stable isotopes of oxygen, carbon and sulfur. *Geochimica Et Cosmochimica Acta* **36**:1337.
- Hobson, K. A. 1999. Tracing origins and migration of wildlife using stable isotopes: a review. *Oecologia* **120**:314-326.
- Hobson, K. A., and K. J. Kardynal. 2016. An isotope (delta S-34) filter and geolocator results constrain a dual feather isoscape (delta H-2, delta C-13) to identify the wintering grounds of North American Barn Swallows. *Auk* **133**:86-98.
- Lott, C. A., T. D. Meehan, and J. A. Heath. 2003. Estimating the latitudinal origins of migratory birds using hydrogen and sulfur stable isotopes in feathers: influence of marine prey base. *Oecologia* **134**:505-510.
- Newsome, S. D., P. W. Collins, T. C. Rick, D. A. Guthrie, J. M. Erlandson, and M. L. Fogel. 2010. Pleistocene to historic shifts in bald eagle diets on the Channel Islands, California. *Proceedings of the National Academy of Sciences of the United States of America* **107**:9246-9251.
- O'Leary, M. H. 1988. Carbon isotopes in photosynthesis. *Bioscience* **38**:328-336.
- Reisinger, A. J., D. T. Chaloner, J. Ruegg, S. D. Tiegs, and G. A. Lamberti. 2013. Effects of spawning Pacific salmon on the isotopic composition of biota differ among southeast Alaska streams. *Freshwater Biology* **58**:938-950.
- Schindler, D. E., M. D. Scheuerell, J. W. Moore, S. M. Gende, T. B. Francis, and W. J. Palen. 2003. Pacific salmon and the ecology of coastal ecosystems. *Frontiers in Ecology and the Environment* **1**:31-37.
- Yerkes, T., K. A. Hobson, L. I. Wassenaar, R. Macleod, and J. M. Coluccy. 2008. Stable isotopes (δD , $\delta^{13}\text{C}$, $\delta^{15}\text{N}$) reveal associations among geographic location and condition of Alaskan northern pintails. *Journal of Wildlife Management* **72**:715-725.