


**Sea Duck Joint Venture**  
**Annual Project Summary for Funded Project**  
**FY 05 – (October 1, 2004 to Sept 30, 2005)**



**Project Title:** No. 4. Atlantic Seaduck Study: Movements, Habitat Use and Feeding Ecology of Seaducks in Chesapeake Bay and Other Atlantic Coastal Areas.

**Principal Investigator(s):** Matthew C. Perry, USGS Patuxent Wildlife Research Center, 12100 Beech Forest Drive, Laurel, MD 20708, USA. [Matt\\_Perry@usgs.gov](mailto:Matt_Perry@usgs.gov)

**Partners:** USFWS and CWS

**Project Description:** The migrational pathways and critical habitats used by seaducks in the Atlantic Flyway are in need of further research, in particular, determination of the breeding and molting areas of surf scoters (*Melanitta perspicillata*) and black scoters (*Melanitta nigra*).

During March-April 2001 - 2004 surf scoters were live-captured in Chesapeake Bay, Maryland, USA. Initial work with floating mist nets and night-lighting was unsuccessful in capturing scoters in Chesapeake Bay. A net gun (Coda, Inc.) fired from a fast-moving boat at approximately 5-10 meters from the boat captured all surf scoters. Standard night-lighting procedures were used on the Restigouche River in New Brunswick to capture black scoters during May 2002 - 2004. Night-lighting techniques use hand nets to capture ducks from the bow of a slowly moving boat with the aid of bright lights powered by batteries or generators.

Males and females were used for the telemetry aspect of this study as both sexes go to the breeding areas, although males leave shortly after initiation of incubation and go to molting areas. Within 24 hours after capture, each scoter underwent an intra-abdominal surgery to implant PTT 100 satellite transmitters (39 g) manufactured by Microwave, Inc., Columbia, Maryland. The transmitter's configuration included an external antenna (percutaneous) that was passed through the back of the duck using a surgical catheter. Initially, each duck was held post-surgery for 3-9 days, but during 2003 and 2004 ducks were released within 24 hours of instrumentation. All ducks were released at the site of capture.

**Objectives:** The study goal was to determine the current distribution of surf and black scoters on their breeding and molting areas. Delineating these populations and establishing the affinities among wintering, staging, breeding, and molting grounds will assist in future population monitoring and management. Specific objectives were:

1. Determine migrational route and breeding ground of surf scoters, when they leave Chesapeake Bay in the late-winter, and black scoters, when they leave Restigouche River in June.
2. Conduct habitat analyses using detailed maps to provide data on the hydrology and geomorphology of these remote scoter breeding areas and the important biotic and abiotic factors influencing nest site selection.
3. Conduct habitat analyses of staging, molting and wintering habitats of black and surf scoters, based on satellite telemetry coordinates.

**Preliminary Results:** A total of 17 surf scoters have been instrumented during four years (2001 - 2004) of satellite telemetry studies and 32 black scoters have been instrumented during three years (2002 - 2004). Males and females have been instrumented during this study. By instrumenting male scoters we have been able to get information of their movements to breeding areas, but then also obtain information about their movements to molting areas. Males only remain on the breeding area for approximately 2-3 weeks, so by instrumenting females we can obtain more information about breeding areas and possible movement with young. Updated information from the ARGOS Systems aboard the NOAA satellites on scoter movements was made accessible on the Patuxent Website: <http://www.pwrc.usgs.gov/resshow/perry/scoters/>.

Satellite telemetry determined that James Bay was the major molting area for male black and surf scoters, although a few males molted along the coast of Labrador-Newfoundland. Black scoters that were instrumented on the Restigouche River, which is a major staging area, were widely distributed along the Atlantic Coast from Cape Cod to Georgia during winter. Major wintering areas for black scoters were Cape Cod (Martha's Vineyard and Nantucket Island), Long Island, and New Jersey.

Only one of the 49 scoters that was instrumented died during migration and this was a male black scoter that was legally shot in December 2002 near Atlantic City, New Jersey. Surf scoters instrumented in Chesapeake Bay in late winter showed a strong tendency to return to Chesapeake Bay for the following winter after they had migrated to and from breeding areas in Quebec.

During this past fiscal year, we analyzed characteristics of wetland habitats used by breeding black scoters and surf scoters in northeastern North America based on satellite telemetry data collected in the spring and summer. During 2002 and 2004, nine black scoters (four males, five females) were tracked to nesting areas in Quebec, Manitoba, and Northwest Territories. In addition, in 2001-04, seven surf scoters (three males, four females) were tracked to nesting areas in Newfoundland and Labrador, Quebec, Northwest Territories, and Nunavut. Based on satellite telemetry data, locations of black and surf scoters in breeding areas on average were not significantly different in regard to latitude and longitude. Presumed nesting sites were manually plotted on the appropriate topographic maps and percent cover type and water were estimated. Nesting habitat of black scoters was significantly different than nesting habitat of surf scoters, with black scoters mainly using open areas (44%) and surf scoters using mainly forested areas (66%). Surf scoters presumed nesting sites were at significantly higher elevations than

sites used by black scoters. Some nesting sites were associated with islands, but the role of islands for nesting sites is equivocal. These results aid in the identification of critical breeding areas and provide a baseline classification of nesting habitats used by these two species.

Water quality and classification data were collected on other nesting lakes in Quebec (Lac Malbaie and Hydro-Quebec areas) and Labrador (Goose Bay area) that were more accessible to researchers. We examined whether a correlation existed among characteristics of lakes used by scoter breeding pairs compared to the characteristics of lakes not used by scoters. The following parameters were measured: dissolved oxygen, carbon dioxide, pH, minerals, nitrate, phosphate, and secchi disk light penetration. Aquatic vegetation, sediment type, benthic invertebrates, size and shape of lake, presence of islands, water sources (marsh, lake, river, etc.) were also determined. Dissolved oxygen and pH were the only parameters that significantly differed among the lakes examined. The mean depth of lakes, the presence of islands, and the presence of predatory fish are potential factors controlling selection of nesting habitat for black and surf scoters in northeastern North America, and more research is necessary to determine the importance of these factors. This information will be useful to resource managers in the future who are responsible for the remote habitats of these scoter species.

During 2004, 10 black scoters and 5 surf scoters were successfully instrumented and tracked from wintering and staging areas to breeding and molting areas. We obtained good movement data from the scoters and several were tracked to new breeding areas in Nunavut and Northwest Territories. The new breeding sites emphasized the fact that the sample size with satellite telemetry is inherently small and several years of data are necessary to obtain desired information. We hope future work with 5 female surf and 5 female black scoters will provide new data about the movements of these scoter species and the habitat they use during migration and on wintering, breeding, and molting areas.

**Project Status:**

During FY 2005 only three surf scoters were instrumented due to lack of funds. Transmitters from these three ducks were donated by Microwave Telemetry, Inc. Tracking of these three ducks is ongoing and all data is presented on our website which is updated weekly. <http://www.pwrc.usgs.gov/resshow/perry/scoters/>